arXiv:hep-ph/9809516v2 12 Oct 1998

Infrared and light-cone limit in hot QED*

E. Petitgirard
Gesellschaft für Schwerionenforschung mbH, Planckstr. 1,
64291 Darmstadt, Germany

In hot gauge theories a breakdown of the hard thermal loop expansion occurs for light-like external momenta or in the infrared region. In QED where a resummation of ladder diagrams is usually advocated, it is shown that long range magnetic interations involve a broader set of graphs. The consequence is a generalized compensation of the hard modes damping terms at leading order in the infrared limit and near the light-cone. The relevance of the so-called improved hard thermal loop resummation scheme is discussed.

I. INTRODUCTION

At the scale of the order of the Debye and plasmon masses, gauge theories at high temperature can no longer be adequately described by ordinary thermal perturbation theory. A consistent approach requires the so-called hard thermal loop expansion [1–6]. Although successful in resolving many paradoxes, there still remain some fundamental problems in certain limits. A particular problem is that of the damping rate of a fast fermion, where self-consistent schemes outside of the hard thermal loop expansion have been used [7,8]. Another class of problems arises whenever soft external momenta are light-like. This is the case with the soft photon production rate estimates [9,10]. Basically, like the hard thermal loops themselves, the polarization tensor involved in the calculation of such processes develops collinear singularities. In the same context, a resummation of asymptotic thermal masses for hard modes, the so-called improved hard thermal loop scheme, leaves the gauge invariance of the effective action intact while screening the divergences [11,12].

But in these cases, another candidate for the removal of the collinear singularities would be the damping term for hard modes. This was suggested for example in another estimate of the soft photon production rate [13]. Indeed, through the interactions with soft modes, the damping is anomalously large $\gamma \sim g^2 T \ln(1/g)$. However if γ has to be taken into account, modifying only the propagators violates the Ward identities. In Abelian theories the latter give $Q^\mu \Pi_{\mu\nu} = 0$, but it turns out that

$$Q^\nu \Pi_{0\nu}(q_0, 0) = \frac{2i\gamma q_0}{q_0 + 2i\gamma} \frac{e^2 T^2}{3} \tag{1}$$

Vertex corrections are necessary to restore gauge invariance and leads to a ladder resummation. The latter arise for instance in the context of the eikonal expansion of gauge theories [14,15]. In the infrared limit of the polarization tensor in hot QED [7], using a constant damping, it was found that the damping term gets eliminated from all the components of $\Pi^{\mu\nu}$. Subsequent works [12,16] in scalar QED have reached the same conclusion for specific limits. Recently a general way to get the cancellation of the damping term, via simple algebraic compensations, has been put forward [17,18]. Besides the non-transversality of the polarization tensor mentioned above, there is another argument concerning gauge invariance if the case of a non-constant damping is considered. It is known that when evaluating the damping rate, keeping the external line on mass-shell with a finite infrared cutoff leads to a gauge independent contribution [19]. But with fermion propagators related to specific internal lines of the polarization tensor, the integration over the real axis usually leaves the momentum off mass-shell. For hard fermions there are gauge dependent pieces off mass-shell besides the factor $e^2 T \ln(1/e)$, at least when going beyond a simple logarithmic approximation. A general compensation of the damping term seems therefore necessary.

These considerations do not concern directly the estimates of the soft photon production rate. The definition of the latter (in covariant gauges) involves the trace Π^μ_μ. This term is subleading compared to the order of the hard thermal loops (HTL) and requires more involved calculations. The approximations justified at the HTL level and which are presented in the following are no longer valid in this case. Nevertheless the soft photon rate calculations share basic features with the current analysis. The fact that ladders contribute, both in the infrared and the light-cone limit, at the same leading order as the HTL is due to a resonance of denominators [7]. The same resonance of denominators occurs for the soft photon rate and leads to a mechanism of enhancement when regularizing the collinear divergences with the fermion asymptotic mass [10]. The bremsstrahlung processes dominate over previous estimates based on the soft fermion loop computations [9]. Now the bremsstrahlung diagrams can be seen as a first step towards a resummation of higher order graphs. This is highly expected, from the considerations above concerning $\Pi^{\mu\nu}$ at leading order on the light-cone, and also from semi-classical calculations taken into account the Landau-Pomeranchuk-Migdal effect [20]. Finally the following study about the cancellation of damping terms casts doubt concerning attempts to regularize the photon rate in the manner proposed in Ref. [13], even if a direct investigation remains to be done.

---

*Talk presented at the 5th International Workshop on Thermal Field Theories and their Applications, Regensburg, Germany, August 1998

Therefore, already at the order of the HTL, an important motivation is to make explicit the equivalence between the infrared and the light-cone limit. In that respect there are two technical new points compared to previous studies about ladders. The first one is that Ref. [17] left ambiguities with an ill-defined expression for the vertex solution of the Bethe-Salpeter equation. Such ambiguities prevented a rigorous cancellation of damping terms. This problem can be overcome by reproducing a structure similar to the Landau damping contributions at the HTL level, *i.e.* a separation between a positive energy and a negative energy part. The second point is that diagrammatically, in order to consider a compensation of a non-constant damping rate, it is necessary to go beyond a simple ladder resummation by taking into account at leading order a more general set of graphs. Thus in the infrared region the usual HTL term will be recovered, as it was already expected from arguments using simplified models (constant damping) and using also the kinetic approach [21,22]. Another motivation is to check the validity of the so-called improved HTL resummation scheme on the light-cone.

II. LEADING GRAPHS

Several works have been devoted to the cancellation of ladder graphs in an effective expansion. It is therefore not necessary to enter into details on this topic. But it is worth recalling, even very briefly, the power counting argument which leads to a ladder resummation. The retarded/ advanced formalism [23] is used, since the structure of Green functions as tree-like diagrams can easily be seen. However all the calculations could be performed using a different real time formalism, for example the Schwinger-Keldysh technique [24]. An important point is that leading terms are given by trees associated with cut internal lines for the photons. For the simple one-loop $ee\gamma$ vertex where Q, P and $R = P + Q$ are respectively the photon momentum and the incoming and outgoing fermion momenta, and the particular example of an exchanged transverse photon of momentum L, splitting the product of fermion propagators $\Delta(P+L)\Delta(R+L)$ along the ways describes in Ref. [17] gives

$$\tilde{V}^{\mu}_{RAR}(P,Q,-R) = -e^2 \int \frac{d^4L}{(2\pi)^4} P^t_{\rho\sigma}(L)\gamma^\rho(\not{R}+\not{L})\gamma^\mu (\not{P}+\not{L})\gamma^\sigma n(l_0)\rho_T(L)\frac{1}{2Q.(P+L)+Q^2}\left(\Delta_A(R+L) -\Delta_R(P+L)\right). \tag{2}$$

An estimate of the expression above can be done if the denominator $2Q.(P+L)+Q^2$ is of the same order as $2P.Q+Q^2$

$$\tilde{V}^{\mu}_{RAR}(P,Q,-R) \sim -e^2\frac{2P^\mu+Q^\mu}{2Q.P+Q^2}\int\frac{d^4L}{(2\pi)^4}P^t_{\rho\sigma}(L) \gamma^\rho \not{P}\gamma^\sigma n(l_0)\rho_T(L)\left(\Delta_A(R+L)-\Delta_R(P+L)\right). \tag{3}$$

Under what circumstances this approximation can actually be made and the denominator extracted from the integral remains to be seen. The difference of the self-energies written above has the same order as the damping rate, namely $O(e^2T)$. This is compensated by the factor $1/e^2$ from the term $1/(2P.Q+Q^2)$. The vertex is therefore of the same order as its tree-level counterpart and this implies a resummation of ladder diagrams



with the exchanges of both transverse and longitudinal soft photons. The reasoning above was made with bare fermion propagators, but when a hard electron or positron is close to its mass-shell, a resummation of its self-energy is necessary. The latter involves one-loop contributions with a bare fermion propagator for the longitudinal photon case and a resummed fermion propagator for the transverse photon exchange.

It can be shown that the ladders cancel against the corresponding self-energies without vertex correction. This is due to a mechanism related to the Ward identities. But with the Landau damping part of the transverse photon spectral density and the absence of Debye screening, the momenta can reach the very soft scale $O(e^2T)$. This may change the estimate of multi-loop diagrams. If self-energies with vertex corrections contribute at leading order as well, as it was pointed out in Ref. [25], then graphs other than ladders might be relevant for consistency and it is necessary to go beyond what has been done in previous works [7,12,17,18]. An estimate of the multi-loop graphs can be made using the simplified transverse spectral density introduced in Ref. [25]

$$\frac{\rho_T(Q)}{q_0} = \frac{2\pi}{q^2}\delta(q_0). \tag{4}$$

An evaluation of the imaginary part of the two-loop self-energy, with P on-shell, yields

$$\frac{1}{4p_0}\mathrm{Tr}\left(\not{P}\Sigma_{RR}(P)\right) = (e^2T)^2\frac{10i}{3(2\pi)^3}\frac{1}{\mu}, \tag{5}$$

where μ is a lower cutoff of order $O(e^2T)$. This result can also be derived within a more complete and rigorous analysis [26] using well known sum rules (see for instance Refs. [27,8]). It justifies the fact that, within a pure logarithmic accuracy, such graphs were discarded [7], but become necessary when going beyond this approximation in order to consider a general (non-constant) damping term.

For consistency this last remark implies that the vertex for a photon in the infrared region or near the light-cone must contain at least, not only ladders, but ladders with vertex corrections. It turns out that non-planar diagrams like the crossed graph



contribute also at the same leading order as the tree-level vertex. The crossed diagram is of the same order as the quantity

$$\tilde V^{+^{\mu}}_{RAR}(P,Q,-R) \sim \frac{\hat P^{\mu}}{q_0+\Omega_P-\Omega_R}(e^2T)^2\frac{2}{(2\pi)^3} \times\frac{1}{\mu}\ln\left(\frac{\omega_p}{\mu}\right), \qquad (6)$$

and is therefore not suppressed by an extra factor e^2 contrary to what has been claimed in Refs. [17,18]. In this estimate μ and the plasma frequency ω_p are respectively the lower (of order $O(e^2T)$) and upper (of order $O(eT)$) cutoffs, Ω_P and Ω_R the moduli of $\vec p$ and $\vec r$.

As for graphs with hard photon exchanges non-trivial compensations between the trace and some denominators kill the mechanism of enhancement encountered before. Therefore only diagrams with soft transverse photons have to be added and their resummation along with the ladder expansion is summarized in the Bethe-Salpeter equation with an infinite kernel. The latter is the sum of one ladder, the subset of 'two-loop' graphs, 'three-loop' graphs etc... Fermion propagators are resummed and, besides the 'one-loop' diagrams mentioned above, a 'two-loop' vertex correction with transverse photons, 'three-loop', etc...must be added. This is consistent with the mechanism of cancellation of self-energies in the polarization tensor which now has to be investigated.

III. RESUMMED DIAGRAMS

A method for including the relevant diagrams when going beyond the logarithmic approximation must now be considered. This is summarized in the Bethe-Salpeter equation

$$\tilde V^{\mu}(P,Q,-R) = \gamma^{\mu} + \int \frac{d^4P'}{(2\pi)^4} K(P,-P',R',-R)\not K_R \not R' \Delta(R')\tilde V^{\mu}(P',Q,-R')\not P'\Delta(P')\not K_P, \qquad (7)$$

where

$$\begin{aligned} K(P,-P',R',-R)\not K_P\cdot\not K_R &= K^1(P,-P',R',-R) \\ &\times\not K^1_P\cdot\not K^1_R + K^2(P,-P',R',-R)\not K^2_P\cdot\not K^2_R+... \\ R=P+Q; &\qquad R'=P'+Q, \end{aligned}$$

which involves an infinite kernel of four-fermion amplitudes that cannot be disconnected by cutting two fermion lines (to avoid overcounting). The first term of this equation represents the simple ladder resummation with resummed fermion propagators and soft longitudinal and transverse photons. The second term contains the summation of the 'two-loop' graphs, namely the crossed diagram and the two ladders with one vertex correction. These diagrams involve only (very) soft transverse photons, as the absence of Debye screening allows to reach the scale $O(e^2T)$ which compensates the extra factor e^2 when adding a further loop. The subsequent contributions describe the resummation of the "three-loop', 'four-loop' graphs, etc...

Now the resummed fermion propagators involves the self-energy at leading order when P^2 is close to the mass-shell. First this leading order contribution includes the *hard thermal loop* term with the effective (and constant) asymptotic mass for a hard fermion

$$\begin{aligned} \Sigma_{HTL}(P) &= ap_0\gamma_0 + b\vec p.\vec\gamma, \\ m^2_\infty &= 2(p_0^2a+p^2b) = \frac{1}{4}e^2T^2. \end{aligned} \qquad (8)$$

Second it must contain also all the exchanges with soft photons. It is then useful to define the quantity

$$\sigma_R(P) = -\frac{1}{4p_0}\mathrm{Tr}\left(\not P\Sigma_R(P)\right), \qquad (9)$$

the imaginary part of which just corresponds to the (general and non-constant) damping rate. The propagator reads

$$\begin{aligned} S_R(P) &= \frac{i}{\not P - \Sigma_{HTL}(P) - \Sigma_R(P)} \\ &= \frac{i\not P + O(e^2)}{P^2 - m^2_\infty + 2p_0\sigma_R(P)}. \end{aligned} \qquad (10)$$

Discarding the damping term would allow to recover the form advocated in ref. [11]. An important starting point is to decompose the resummed propagator into a positive energy part and its negative counterpart

$$\begin{aligned} \Delta_{R/A}(P) &= \frac{i}{P^2 - m^2_\infty + 2p_0\sigma_{R/A}(P)} \\ &\simeq \frac{i}{2p}\left(\frac{1}{p_0-\Omega_P+\sigma^+_{R/A}(P)} - \frac{1}{p_0+\Omega_P+\sigma^-_{R/A}(P)}\right) \\ &= \Delta^+_{R/A}(P) + \Delta^-_{R/A}(P), \end{aligned} \qquad (11)$$

where $\Omega_P = [\vec{p}^2 + m_\infty^2]^{1/2}$. Keeping in mind this canonical decomposition, an estimate [26] enables to guess the expression of an *improved* vertex

$$\tilde{V}^\mu_{RAR}(P,Q,-R) = \gamma^\mu + \frac{v^\mu}{q_0 + \Omega_P - \Omega_R}\left(\Sigma_R^+(P)\right.$$
$$\left.-\Sigma_A^+(R)\right) + \frac{\bar{v}^\mu}{q_0 + \Omega_R - \Omega_P}\left(\Sigma_R^-(P) - \Sigma_A^-(R)\right), \quad (12)$$

where new vectors can be defined

$$v^\mu = \left(1, \left(1 - \frac{m_\infty^2}{2p^2}\right)\hat{p}^i\right),$$
$$\bar{v}^\mu = \left(1, -\left(1 - \frac{m_\infty^2}{2p^2}\right)\hat{p}^i\right), \quad (13)$$

in order for the above vertex to satisfy Ward identities. These vectors are equal to the unit vectors $(1, \hat{p}^i)$ and $(1, -\hat{p}^i)$ plus soft contributions unimportant in the determination of the leading order part of the polarization tensor.

It is now necessary to check that for very soft external photons, this vertex is indeed the solution of the Bethe-Salpeter solution, even though more complicated diagrams than the ladders have been included. It is also important to see to what extent it is still a solution of (7) in the light-cone limit.

IV. LADDER RESUMMATION

The simple ladder resummation has been extensively considered in the litterature. Here the additional input is the separation of the expression of the vertex between a positive energy part and its negative counterpart. This results from the canonical decomposition of the resummed fermion propagators and due to the basic assumption of an energetic fermion emitting or absorbing soft photons. Sticking the expression (12) into the first term of the Bethe-Salpeter equation yields

$$\tilde{V}^{1^\mu}_{RAR}(P,Q,-R) = \sum_{i=t,l}(-e^2)\int [dL]^i_{\rho\sigma}\Delta_R(P+L)$$
$$\Delta_A(R+L)\gamma^\rho(\not{R}+\not{L})\tilde{V}^\mu_{RAR}(P+L,Q,-R-L)$$
$$(\not{P}+\not{L})\gamma^\sigma, \quad (14)$$

where the notation $[dL]$ for either a transverse or a longitudinal spectral density stands for

$$[dL]^i_{\rho\sigma} = \frac{d^4L}{(2\pi)^4} n(l_0)\rho_i(L) P^i_{\rho\sigma}(L). \quad (15)$$

Contracting the spinors gives

$$(\not{R}+\not{L})\gamma^\mu(\not{P}+\not{L}) \simeq 2P^\mu \not{P},$$
$$(\not{R}+\not{L})v^\mu\Sigma^+(P+L)(\not{P}+\not{L}) \simeq -2P^\mu\sigma^+(P)\not{P}. \quad (16)$$

Soft terms may be discarded and the basic picture of an energetic electron (or positron) undergoing interactions with soft photons (momenta of order $O(eT)$ at most) implies that the former is always close to its mass-shell ($P^2 \sim O(eT^2)$ at most). This allows to make several approximations in the equations above. The expression (14) now reads

$$\tilde{V}^{1^\mu}_{RAR}(P,Q,-R) = \sum_{i=t,l}(-e^2)\int [dL]^i_{\rho\sigma}\Delta_R(P+L)$$
$$\Delta_A(R+L)2P^\mu\gamma^\rho\not{P}\gamma^\sigma\left[1 - \frac{\sigma_R^+(P+L) - \sigma_A^+(R+L)}{q_0 + \Omega_{P+L} - \Omega_{R+L}}\right.$$
$$\left.-\frac{\sigma_R^-(P+L) - \sigma_A^-(R+L)}{q_0 + \Omega_{R+L} - \Omega_{P+L}}\right]. \quad (17)$$

This enables also to split the product of propagators $\Delta(P)\Delta(R)$ according to their canonical decompositions (mixed positive/ negative energy terms are shown to be subleading)

$$\Delta_R(P)\Delta_A(R) \simeq \frac{i}{2p}\left(\frac{\Delta_R^+(P) - \Delta_A^+(R)}{q_0 + \Omega_P - \Omega_R - \sigma_R^+(P) + \sigma_A^+(R)}\right.$$
$$\left.-\frac{\Delta_R^-(P) - \Delta_A^-(R)}{q_0 + \Omega_R - \Omega_P + \sigma_A^-(R) - \sigma_R^-(P)}\right), \quad (18)$$

and to relate the positive or negative energy parts of the propagators to the corresponding contributions of the vertex

$$\left[\Delta_R^+(P) - \Delta_A^+(R)\right]\left[1 - \frac{\sigma_R^+(P) - \sigma_A^+(R)}{q_0 + \Omega_P - \Omega_R}\right.$$
$$\left.-\frac{\sigma_R^-(P) - \sigma_A^-(R)}{q_0 + \Omega_R - \Omega_P}\right]$$
$$\simeq \left[\Delta_R^+(P) - \Delta_A^+(R)\right]\left[1 - \frac{\sigma_R^+(P) - \sigma_A^+(R)}{q_0 + \Omega_P - \Omega_R}\right]. \quad (19)$$

It can be easily seen that the algebraic cancellation occurs quite naturally using this decomposition. The first term of the Bethe-Salpeter equation equation may now be written

$$\tilde{V}^{1^\mu}_{RAR}(P,Q,-R) = \sum_{i=t,l}(-ie^2)\int [dL]^i_{\rho\sigma}\gamma^\rho\not{P}\gamma^\sigma$$
$$\left(v^\mu\frac{\Delta_R^+(P+L) - \Delta_A^+(R+L)}{q_0 + \Omega_{P+L} - \Omega_{R+L}}\right.$$
$$\left.+\bar{v}^\mu\frac{\Delta_R^-(P+L) - \Delta_A^-(R+L)}{q_0 + \Omega_{R+L} - \Omega_{P+L}}\right). \quad (20)$$

It is now crucial to see to what extent the denominators without damping can be extracted from the integral. Expanding their expressions with the soft terms from L and Q gives for example

$$q_0 + \Omega_{P+L} - \Omega_{R+L} = q_0 + \Omega_P - \Omega_R$$
$$-\frac{\vec{q}}{p}.\left[\vec{l} - \hat{p}(\hat{p}.\vec{l})\right] + O(e^3T). \quad (21)$$

In the infrared limit ($q_0, q \sim O(e^2T)$) and on the light-cone for a not too collinear configuration ($\hat{p}.\hat{q} \sim \pm 1 + O(e)$) the terms following $q_0 + \Omega_P - \Omega_R$ may always be neglected. This is obvious in the infrared region. For the light-cone limit it turns out that $\hat{q} \sim \hat{p}$ and consequently $(\vec{q}/p).(\vec{l} - \hat{p}(\hat{p}.\vec{l})) \sim O(e^3T)$. However one of the main points is to notice that in a more collinear configuration these subsequent contributions are no longer neglegible. Indeed in this last case $q_0 + \Omega_P - \Omega_R$ is of the same order $O(e^3T)$. This is the situation where the effect of the asymptotic mass is important. Therefore, outside this limit, *i.e.* in the infrared and in a 'weak' light-cone limit, the relevant denominators may be extracted and the expression of the vertex finally gives

$$\tilde{V}^{1^\mu}_{RAR}(P,Q,-R) = \frac{v^\mu}{q_0 + \Omega_P - \Omega_R}\left(\Sigma^{1^+}_R(P) - \Sigma^{1^+}_A(R)\right) + \frac{\bar{v}^\mu}{q_0 + \Omega_R - \Omega_P}\left(\Sigma^{1^-}_R(P) - \Sigma^{1^-}_A(R)\right). \tag{22}$$

But of course this part does not correspond to the the full contribution of the vertex (12) since it is necessary to go beyond a simple ladder resummation.

V. N-LOOP DIAGRAMS RESUMMATION

So far only a simple ladder expansion has been considered. Since also 'higher-order' graphs have been shown to contribute at leading order it is therefore necessary to take into account their resummation. This corresponds to an infinite kernel for the Bethe-Salpeter equation.

As a first step beyond the ladders, the algebraic cancellation of self-energies for the 'two-loop' graphs must now be considered. Inserting the vertex (12) gives for the 'two-loop' crossed diagram

$$\begin{aligned}\tilde{V}^{2^\mu}_{Cr_{RAR}}(P,Q,-R) &= e^4\int [dL_1]^t_{\alpha\beta}[dL_2]^t_{\sigma\rho}\Delta_A(R+L_1)\\ &\Delta_A(R+L_1+L_2)\Delta_R(P+L_1+L_2)\Delta_R(P+L_2)Sp(R)\\ &\tilde{V}^\mu_{RAR}(P+L_1+L_2,Q,-R-L_1-L_2)Sp(P),\end{aligned} \tag{23}$$

where the Δ's are the resummed fermion propagators and

$$\begin{aligned}Sp(R) &= \gamma^\alpha(\not{R}+\not{L}_1)\gamma^\sigma(\not{R}+\not{L}_1+\not{L}_2),\\ Sp(P) &= (\not{P}+\not{L}_1+\not{L}_2)\gamma^\beta(\not{P}+\not{L}_2)\gamma^\rho.\end{aligned} \tag{24}$$

The product of propagators $\Delta(P+L_1+L_2)\Delta(R+L_1+L_2)$ attached to the internal vertex has to be split in the same way as for the ladders. The decomposition of fermion propagators into positive and negative energy parts allows to separate a positive energy contribution from its negative counterpart in the vertex. The algebraic cancellation still occurs and is expressed through the replacements of $(q_0 \pm \Omega_{P+L_1+L_2} \mp \Omega_{R+L_1+L_2} \pm \sigma_{R_{P+L_1+L_2}} \mp \sigma_{A_{R+L_1+L_2}})$ by the denominators $(q_0 \pm \Omega_{P+L_1+L_2} \mp \Omega_{R+L_1+L_2})$. Finally in the infrared and in a weak light-cone limit the L_i-dependent terms can always be neglected and $(q_0+\Omega_P-\Omega_R)$ or $(q_0+\Omega_R-\Omega_P)$ turns out to be a good aproximation for $(q_0+\Omega_{P+L_1+L_2}-\Omega_{R+L_1+L_2})$ or $(q_0+\Omega_{R+L_1+L_2}-\Omega_{P+L_1+L_2})$. Extracting the first denominators from the integral yields

$$\begin{aligned}\tilde{V}^{2^\mu}_{Cr_{RAR}}(P,Q,-R) &= \frac{v^\mu}{q_0+\Omega_P-\Omega_R} ie^4\int [dL_1]^t_{\alpha\beta}[dL_2]^t_{\sigma\rho}\\ &Sp(P,R)\Delta^+_A(R+L_1)\Delta^+_R(P+L_2)\left(\Delta^+_R(P+L_1+L_2)\right.\\ &\left.-\Delta^+_A(R+L_1+L_2)\right) + n.e.\end{aligned} \tag{25}$$

with

$$Sp(P,R) = \gamma^\alpha(\not{R}+\not{L}_1)\gamma^\sigma(\not{P}+\not{L}_1+\not{L}_2)\gamma^\beta(\not{P}+\not{L}_2)\gamma^\rho, \tag{26}$$

(the terms L_i in the numerator are subleading but have been kept to stress the similarity with Ward identities). If now the other 'two-loop' diagrams are considered, namely the vertex correction attached to the soft photon line L_1 and its symmetric counterpart, the same procedure as before can be applied when inserting the full vertex. In particular the resulting denominators $(q_0 \pm \Omega_{P+L_i} \mp \Omega_{R+L_i})$ may be approximated by $(q_0 \pm \Omega_P \mp \Omega_R)$ in the aforementioned limits. Adding the positive energy and negative energy contributions of these diagrams with the crossed graph gives

$$\tilde{V}^{2^\mu}_{RAR}(P,Q,-R) = \frac{v^\mu}{q_0 + \Omega_P - \Omega_R}\left(\Sigma^{2^+}_R(P) - \Sigma^{2^+}_A(R)\right) + \frac{\bar{v}^\mu}{q_0 + \Omega_R - \Omega_P}\left(\Sigma^{2^-}_R(P) - \Sigma^{2^-}_A(R)\right). \tag{27}$$

The reasoning above can be generalized to the 'N-loop' vertices in the Bethe-Salpeter equation. These vertices involve resummed fermion propagators and the exchange of soft transverse photons. They induce four-fermion amplitudes which cannot be disconnected by cutting two fermion lines. Each of these graphs have internal legs $P+L_i+...+L_j$ and $R+L_i+...+L_j$ attached to the external photon line via the complete vertex. Splitting the product of propagators $\Delta(P+L_i+...+L_j)\Delta(R+L_i+...+L_j)$ as before leads to the algebraic cancellation of the self-energies σ, *i.e.* to the replacement of the common denominator $(q_0+\Omega_{P+L_i+...}-\Omega_{R+L_i...}+\sigma_{R_{P+L_i+...}}-\sigma_{A_{R+L_i+...}})$ by $(q_0+\Omega_{P+L_i+...}-\Omega_{R+L_i...})$. Again in the infrared region and near the light-cone without a too strong collinearity between the photon and the electron the latter denominator can be approximated by $q_0+\Omega_P-\Omega_R$ ($q_0+\Omega_R-\Omega_P$ for the negative energy part) and extracted from the integral. What is left is completely equivalent to a contraction of each graph with Q^μ, times the vector $v^\mu/(q_0+\Omega_P-\Omega_R)$ (and its negative energy counterpart $\bar{v}^\mu/(q_0+\Omega_R-\Omega_P)$). It is therefore natural for a demonstration at any order to rely on Ward identities between vertices with resummed fermions and cut soft transverse photons and the corresponding self-energies. The other additional input being the separation between a positive energy contribution and the negative one. This is due to the basic assumption of an energetic (hard) fermion interacting with soft photons. Each diagram is related to specific terms which would be obtained with a simple contraction with Q^μ

$$\tilde{V}^{N^\mu}_{M\,K_{RAR}}(P,Q,-R) = \frac{v^\mu}{q_0+\Omega_P-\Omega_R}\left(\Sigma^{N^+}_{M\,K_R}(P) - \Sigma^{N^+}_{M\,K_A}(R)\right) + \frac{\bar{v}^\mu}{q_0+\Omega_R-\Omega_P}\left(\Sigma^{N^-}_{M\,K_R}(P) - \Sigma^{N^-}_{M\,K_A}(R)\right), \tag{28}$$

the sum of which corresponds to a self-energy with a particular vertex correction

$$\Sigma^{N^\pm}_{M_R}(P) = \sum_K \Sigma^{\pm}_{M\,K_R}(P). \tag{29}$$

Adding these self-energies yields the 'N-loop' order Σ

$$\Sigma^{N^\pm}_R(P) = \sum_M \Sigma^{\pm}_{M_R}(P), \tag{30}$$

and finally the contribution of all 'N-loop' self-energies gives the complete leading order expression

$$\Sigma^{\pm}_R(P) = \sum_M \Sigma^{N^\pm}_R(P), \tag{31}$$

which allows to recover the *improved* vertex (12). The latter is therefore the solution of the Bethe-Salpeter equation for the particular cases considered so far.

VI. CANCELLATION OF DAMPING TERMS

In the infrared limit the improved vertex

$$\tilde{V}^\mu_{RAR}(P,Q,-R) = \gamma^\mu + \frac{v^\mu}{q_0+\Omega_P-\Omega_R}\left(\Sigma^+_R(P) - \Sigma^+_A(R)\right) + \frac{\bar{v}^\mu}{q_0+\Omega_R-\Omega_P}\left(\Sigma^-_R(P)-\Sigma^-_A(R)\right), \tag{32}$$

is therefore shown to be the solution of the Bethe-Salpeter equation. In the light-cone limit where the effects of the asymptotic mass are relevant, this is not true any more, since approximations valid in the former case are no longer legitimate. Nevertheless it is always interesting to look at the expression of $\Pi^{\mu\nu}(Q)$ provided by this vertex. To deal with a general (non-constant) damping term, the trick consists in coming back to the imaginary time formalism, comparisons between quantities like $i\omega_n$, Ω_P and $\sigma(i\omega_n,p)$ being easier. In the R/A formalism, the retarded part of the polarization tensor can be written as

$$\begin{aligned} i\Pi^{\mu\nu}_{RR}(Q) = -e^2\mathrm{Tr}\int\frac{d^4P}{(2\pi)^4}\left(\not{P}\gamma^\mu\not{R}\right)\Bigg\{&\left(\frac{1}{2}-n_F(p_0)\right)\\ \Delta_R(R)\left[\tilde{V}^\nu_{RRA}\Delta_R(P) - \tilde{V}^\nu_{ARA}\Delta_A(P)\right] + &\left(\frac{1}{2} - n_F(r_0)\right)\Delta_A(P)\left[\tilde{V}^\nu_{ARA}\Delta_R(R) - \tilde{V}^\nu_{ARR}\Delta_A(R)\right]\Bigg\},\\ \tilde{V}^\nu_{\alpha\beta\delta} = \gamma^\nu + \tilde{V}^{1^\nu}_{\alpha\beta\delta}(P,Q,-R).& \end{aligned} \tag{33}$$

Contracting the spinors gives a part sensitive to the infrared or light-cone limit plus possible tadpole terms. The latter do not involve vertex and damping corrections, or more precisely it can be shown that such corrections are subleading. Splitting the products of propagators yields denominators containing the retarded and advanced damping terms. These denominators cancel against the numerators of the form $1+(\sigma_P-\sigma_R)/(q_0+\Omega_P-\Omega_R)$ derived from the equation (12). Taking Π^{00} as a particular example, a simpler expression is thus obtained

$$\begin{aligned} i\Pi^{00}_{RR}(Q) = -4ie^2\int\frac{d^4P}{(2\pi)^4}p\Bigg\{&\left(\frac{1}{2}-n_F(p_0)\right)\times\\ &\left[\frac{\delta^+(P)}{q_0+\Omega_P-\Omega_R} - \frac{\delta^-(P)}{q_0+\Omega_R-\Omega_P}\right] - \left(\frac{1}{2}-n_F(r_0)\right)\\ &\left[\frac{\delta^+(R)}{q_0+\Omega_P-\Omega_R} - \frac{\delta^-(R)}{q_0+\Omega_R-\Omega_P}\right]\Bigg\},\\ \delta^\pm(P) = \Delta^\pm_R(P) - \Delta^\pm_A(P).& \end{aligned} \tag{34}$$

The integration over p_0 can be converted into a contour integral in the complex half-plane where there are no discontinuities from the propagators but only the poles given by the Matsubara frequencies. The contour can be composed of the real axis, two quarter-circles expanding at infinity, the contribution of which tends to zero, and a part circling the upper or lower poles of the Fermi-Dirac factor. The contribution is finally reduced to the sum of the residues corresponding to the Matsubara frequencies

$$\begin{aligned} i\Pi^{00}_{RR}(Q) = 4i\pi Te^2\int\frac{d^3p}{(2\pi)^3}\sum_n\Bigg[&\frac{1}{q_0+\Omega_P-\Omega_R}\times\\ &\left(\frac{1}{\omega_n-\Omega_P+\sigma^+(\omega_n,p)} - \frac{1}{\omega_n-\Omega_R+\sigma^+(\omega_n,r)}\right)\\ &-\frac{1}{q_0+\Omega_R-\Omega_P}\left(\frac{1}{\omega_n+\Omega_P+\sigma^-(\omega_n,p)}\right.\\ &\left.-\frac{1}{\omega_n+\Omega_R+\sigma^-(\omega_n,r)}\right)\Bigg]. \end{aligned} \tag{35}$$

It is mainly the difference $\Omega_P - \Omega_R \sim \hat{p}.\vec{q}$ which matters. Its order of magnitude is $O(eT)$ in the light-cone limit, and $O(e^2T)$ or below in the infrared region. It is much larger than $\sigma(i\omega_n,p) - \sigma(i\omega_n,r)$ in each case. The contribution due to the damping terms is therefore negligible. The expression given by simple poles in real time is re-established and the improved HTL

$$\begin{aligned} i\Pi^{00}_{RR}(Q) = -4i\pi e^2\int\frac{d^3P}{(2\pi)^4}&\left(n_F(\Omega_R)-n_F(\Omega_P)\right)\\ &\left[\frac{1}{q_0+\Omega_P-\Omega_R}+\frac{1}{q_0+\Omega_R-\Omega_P}\right], \end{aligned} \tag{36}$$

is recovered. The complex energy $q_0 = \mathrm{Re}q_0 + i\epsilon$ allows to get the usual Landau damping part.

In the infrared limit and on the light-cone when $\hat{p}.\hat{q} \sim \pm 1 + O(e)$ the resummation of all the aforementioned vertex diagrams cancels against the self-energy insertions. The result is found to be the *improved* HTL, which just corresponds to the usual HTL in these cases.

However in a strong light-cone limit where the effects of the asymptotic mass are important, the *improved* vertex (12) is no longer solution of the Bethe-Salpeter equation. The expansion (21) is not reduced any more to the first term as it is in the first two cases. The resummation scheme advocated in Ref. [11] is therefore not correct, or at least not complete. However the cancellation of (12) with self-energies whatever the gauge is, provides a further and deeper justification of the gauge invariance of the *improved* HTL of Ref. [11]. But to propose a more complete resummation scheme on the light-cone, other ways of dealing with the Bethe-Salpeter equation are necessary [28].

---

[1] R. D. Pisarski, Physica **A158**, 146, 246 (1989); Phys. Rev. Lett. **63**, 1129 (1989).

[2] V. V. Klimov, Sov. Phys. JETP **55**, 199 (1982).

[3] H. A. Weldon, Phys. Rev. **D26**, 1384, 2789 (1982).

[4] E. Braaten and R. D. Pisarski, Nucl. Phys. **B337**, 569 (1990); **B339**, 310 (1990).

[5] J. C. Taylor and S. M. H. Wong, Nucl. Phys. **B346**, 115 (1990).

[6] J. Frenkel and J. C. Taylor, Nucl. Phys. **B334**, 199 (1990); Z. Phys. **C49**, 515 (1991).

[7] V. V. Lebedev and A. V. Smilga, Ann. Phys. (N.Y.) **202**, 229 (1990); Physica **A181**, 187 (1992).

[8] R. D. Pisarski, Phys. Rev. **D47**, 5589 (1993).

[9] R. Baier, S. Peigné and D. Schiff, Z. Phys. **C62**, 337 (1994); P. Aurenche, T. Becherrawy and E. Petitgirard, *preprint ENSLAPP-A-452-93* (hep-ph/9403320).

[10] P. Aurenche, F. Gelis, R. Kobes and E. Petitgirard, Phys. Rev. **D54**, 5274 (1996); Z. Phys. **C75**, 315 (1997).

[11] F. Flechsig and A. Rebhan, Nucl. Phys. **B464**, 279 (1996).

[12] U. Kraemmer, A. K. Rebhan, and H. Schulz, Ann. Phys. (N.Y.) **238**, 286 (1995).

[13] A. Niégawa, Phys. Rev. **D56**, 1073 (1997); Phys. Rev. **D55**, 4997 (1997).

[14] J. M. Cornwall and G. Tiktopoulos, Phys. Rev. **D15**, 2937 (1977).

[15] J. M. Cornwall and W.-S. Hou, Phys. Lett. **B153**, 173 (1985).

[16] M. Carrington, Phys. Rev. **D48**, 3836 (1993).

[17] M. Carrington, R. Kobes and E. Petitgirard, Phys. Rev. **D57**, 2631 (1998).

[18] M. Carrington and R. Kobes, Phys. Rev. **D57**, 6372 (1998).

[19] A. K. Rebhan, Phys. Rev. **D46**, 4779 (1992).

[20] J. Cleymans, V. V. Goloviznin and K. Redlich, Phys. Rev. **D47**, 989 (1993); Z. Phys. **C59**, 495 (1993).

[21] S. Jeon, Phys. Rev. **D52**, 3591 (1995); S. Jeon and L. Yaffe, Phys. Rev. **D53**, 5799 (1996).

[22] P. Huet and D. T. Son, Phys. Lett. **B393**, 94 (1997).

[23] P. Aurenche and T. Becherrawy, Nucl. Phys. **B379**, 259 (1992); M .A. van Eijck and C. G. van Weert, Phys. Lett. **B278**, 305 (1992).

[24] L. V. Keldysh, Sov. Phys. JETP **20**, 1018 (1964); See also E. M. Lifshitz and L. P. Pitaevsky, *Physical Kinetics* (Pergamon Press, Oxford, 1981).

[25] J. P. Blaizot and E. Iancu, Phys. Rev. **D55**, 973 (1997); Phys. Rev. **D56**, 7877 (1997).

[26] E. Petitgirard, *preprint GSI-98-45* (hep-ph/9808344), to appear in Phys. Rev. **D**.

[27] E. Braaten and T. C. Yuan, Phys. Rev. Lett. **66**, 2183 (1991).

[28] E. Petitgirard, in preparation.